

# Inseparable Orthogonal Matrices over $\mathbb{Z}_2$

Guo-Hui Zhang

*Department of Mathematical Sciences, University of Alabama in Huntsville,  
Huntsville, Alabama 35899*

*Communicated by the Managing Editors*

Received July 29, 1996

A  $(0, 1)$ -matrix  $A$  is called orthogonal over  $\mathbb{Z}_2$  if both  $AA^T$  and  $A^TA$  are diagonal matrices. A matrix  $A$  is called inseparable if  $A$  contains no zero row or zero column and there do not exist permutation matrices  $P$  and  $Q$  such that

[View metadata, citation and similar papers at core.ac.uk](#)

A matrix  $A$  is said to be of type 0 if  $AA^T = O$  and  $A^TA = O$ . A square matrix  $A$  of order  $n$  is said to be of type 1 if  $AA^T = I_n$ . It turns out that an inseparable orthogonal matrix over  $\mathbb{Z}_2$  is either of type 0 or of type 1. Let  $f_0(m, n)$  (respectively,  $F_0(m, n)$ ) denote the smallest (respectively, largest) number of 1's in an  $m \times n$  inseparable orthogonal matrix of type 0 over  $\mathbb{Z}_2$ , and  $f_1(n)$  (respectively,  $F_1(n)$ ) denote the smallest (respectively, largest) number of 1's in an  $n \times n$  inseparable orthogonal matrix of type 1 over  $\mathbb{Z}_2$ . The formulas for  $f_0(m, n)$ ,  $F_0(m, n)$ ,  $f_1(n)$ , and  $F_1(n)$  are completely determined in this paper. © 1998 Academic Press

## 1. INTRODUCTION

In this paper we consider matrices over a given field  $F$ . A *line* of a matrix  $A$  is either a row or a column of  $A$ . Given any  $m \times n$  matrix  $A = (a_{ij})$ , we use  $R_i = R_i(A)$  to denote the  $i$ th row of  $A$  for  $1 \leq i \leq m$ , and  $C_j = C_j(A)$  to denote the  $j$ th column of  $A$  for  $1 \leq j \leq n$ . We also define  $\Omega(R_i) = \{C_j \mid a_{ij} \neq 0\}$  for each  $1 \leq i \leq m$  and  $\Omega(C_j) = \{R_i \mid a_{ij} \neq 0\}$  for each  $1 \leq j \leq n$ . We say a column  $C$  (respectively, a row  $R$ ) *covers* a row  $R$  (respectively, a column  $C$ ) if  $R \in \Omega(C)$ . We define  $\omega(L_1, L_2) = |\Omega(L_1) \cap \Omega(L_2)|$ , where  $L_1$  and  $L_2$  are either two columns or two rows in a matrix  $A$ . The *weight* of a line  $L$ ,  $\omega(L)$ , is the total number of nonzero entries in  $L$ . We also define  $\tau(C) = \sum_{R \in \Omega(C)} \omega(R)$  for any column  $C$  and  $\tau(R) = \sum_{C \in \Omega(R)} \omega(C)$  for any row  $R$ . The *weight* of a matrix  $A$ ,  $\omega(A)$ , is the total number of nonzero entries in  $A$ .

We use  $O$  to denote a zero matrix,  $I_n$  to denote the identity matrix of order  $n$ , and  $J_{m,n}$  to denote the  $m \times n$  all 1 matrix. We abbreviate  $J_{n,n}$  as

$J_n$ . A matrix  $A$  is called *orthogonal* if both  $AA^T$  and  $A^T A$  are diagonal matrices. A matrix  $A$  is said to be of *type 0* if  $AA^T = O$  and  $A^T A = O$ . A matrix  $A$  of order  $n$  over  $F$  is said to be of *type 1* if  $AA^T = A^T A = cI_n$  for some nonzero constant  $c$  in  $F$ . Matrices  $A$  and  $B$  are said to be *p-equivalent* if there exist permutation matrices  $P$  and  $Q$  such that  $PAQ = B$ . A matrix  $A$  of order  $n$  is called *fully indecomposable* if  $A$  is not *p-equivalent* to a matrix of the form  $\begin{pmatrix} B & \\ O & C \end{pmatrix}$ , where  $B$  is a matrix of order  $k$ ,  $1 \leq k \leq n-1$ . A matrix  $A$  is called *admissible* if  $A$  contains no zero row nor zero column. A matrix  $A$  is called *inseparable* if  $A$  is admissible and  $A$  is not *p-equivalent* to a matrix of the form  $\begin{pmatrix} B & O \\ O & C \end{pmatrix}$ , and *separable* otherwise. Clearly, a fully indecomposable matrix is necessarily inseparable. We first state the following result from [5].

LEMMA 1.1. *Let  $A$  be a fully indecomposable, orthogonal matrix over a field  $F$ . Then  $A$  is either of type 0 or of type 1.*

LEMMA 1.2. *Let  $A$  be an inseparable orthogonal matrix over a field  $F$ . If  $A$  is not of type 0, then  $A$  is fully indecomposable and of type 1.*

*Proof.* Let  $A$  be an  $m \times n$  inseparable orthogonal matrix over  $F$ . Assume

$$AA^T = \begin{pmatrix} D_1 & O \\ O & O \end{pmatrix}, \quad A^T A = \begin{pmatrix} D_2 & O \\ O & O \end{pmatrix}, \quad \text{and} \quad A = \begin{pmatrix} A_1 & A_2 \\ A_3 & A_4 \end{pmatrix},$$

where  $D_1$  (respectively,  $D_2$ ) is a diagonal matrix of order  $s$  (respectively,  $t$ ) such that each diagonal entry of  $D_1$  (respectively,  $D_2$ ) is nonzero, and  $A_1$  is the upper-left  $s \times t$  submatrix of  $A$ . Then we have

$$AA^T A = A \begin{pmatrix} D_2 & O \\ O & O \end{pmatrix} = \begin{pmatrix} A_1 D_2 & O \\ A_3 D_2 & O \end{pmatrix}$$

and

$$AA^T A = \begin{pmatrix} D_1 & O \\ O & O \end{pmatrix} A = \begin{pmatrix} D_1 A_1 & D_1 A_2 \\ O & O \end{pmatrix},$$

which implies that  $A_2 = O$  and  $A_3 = O$ . Therefore,

$$A = \begin{pmatrix} A_1 & O \\ O & A_4 \end{pmatrix}.$$

Since  $A$  is inseparable and  $A$  is not of type 0, it follows that  $s = m$  and  $t = n$ . Therefore, we have  $AA^T = D_1$  and  $A^T A = D_2$ . Since the  $m$  rows of  $A$  are linearly independent, it follows that  $m \leq n$ . Similarly, we have  $n \leq m$  since the  $n$  columns of  $A$  are also linearly independent. Therefore, we have  $m = n$  and  $A$  is an invertible matrix of order  $n$ .

We now claim that  $A$  is fully indecomposable. Suppose not, then there exists a submatrix  $B$  of order  $k$ ,  $1 \leq k \leq n-1$ , such that  $A$  is  $p$ -equivalent to a matrix of the form  $\begin{pmatrix} B & C \\ O & D \end{pmatrix}$ . Without loss of generality, we assume  $A = \begin{pmatrix} B & C \\ O & D \end{pmatrix}$ . Then we see that  $C \neq O$  and  $D$  is invertible. Since

$$AA^T = \begin{pmatrix} BB^T + CC^T & CD^T \\ DC^T & DD^T \end{pmatrix}$$

is a diagonal matrix, it follows that  $DC^T = O$ . This would imply that  $C = O$ , which is impossible. Therefore,  $A$  is fully indecomposable, and the result follows immediately from Lemma 1.1. ■

Two vectors  $x = (x_1, \dots, x_n)$  and  $y = (y_1, \dots, y_n)$  are said to be *combinatorially orthogonal* if  $|\{i : x_i y_i \neq 0\}| \neq 1$ . A matrix  $A$  is *combinatorially orthogonal* if each pair of rows of  $A$  is combinatorially orthogonal and each pair of columns of  $A$  is combinatorially orthogonal. Clearly, if  $A$  is an orthogonal matrix over  $F$ , then  $A$  is also combinatorially orthogonal. Furthermore, for any combinatorially orthogonal matrix  $A$  over  $F$ , the matrix obtained from  $A$  by replacing each nonzero entry in  $A$  by a 1 is combinatorially orthogonal. We will need the following result of Reid and Thomassen [4]. The reader is referred to [3] for a simple proof of Lemma 1.3 and further results concerning inseparable and combinatorially orthogonal  $(0, 1)$ -matrices.

LEMMA 1.3. *Let  $A$  be an  $m \times n$  inseparable and combinatorially orthogonal  $(0, 1)$ -matrix. Then  $\omega(A) \geq 2(m+n) - 4$ .*

Let  $f(F; m, n)$  denote the smallest number of nonzero entries in an  $m \times n$  inseparable orthogonal matrix over  $F$ . Then Lemma 1.3 implies that  $f(F; m, n) \geq 2(m+n) - 4$  for any field  $F$ , if it exists. Trivially, we have  $f(F; 2, 2) = 4$  for any field  $F$ . It was proved in [1] that  $f(\mathbb{R}; n, n) = 4(n-1)$ , where  $\mathbb{R}$  is the field of real numbers. It can be shown that  $f(\mathbb{Z}_2; 3, 3)$  does not exist and  $f(F; 3, 3) \geq 8$  if  $F \neq \mathbb{Z}_2$ , with equality if and only if there exists two nonzero elements  $a$  and  $b$  in  $F$  such that  $b^2 - a^2 = 1$ , in which case,

$$A = \begin{pmatrix} a & 1 & a^{-1}b \\ 1 & a^{-1} & -b \\ a^{-1}b & -b & 0 \end{pmatrix}$$

is an orthogonal matrix over  $F$ .

Orthogonal  $(0, 1, -1)$ -matrices over  $\mathbb{R}$  have been referred to as weighing matrices in the literature. Clearly, every weighing matrix can be viewed as an orthogonal matrix over any field  $F$ . Weighing matrices have been

studied extensively (e.g., see [2]). It turns out that the problem of evaluating  $f(F; m, n)$  over a finite field  $F = GF(q)$  is very difficult. For the remainder of this paper, we will consider the case  $F = \mathbb{Z}_2$  only. Note that if a matrix  $A$  of order  $n$  over  $\mathbb{Z}_2$  is of type 1, then we have  $AA^T = A^T A = I_n$ . We also note that if matrices  $A$  and  $B$  are  $p$ -equivalent, then  $\omega(A) = \omega(B)$ , and  $A$  is inseparable (resp., orthogonal) if and only if  $B$  is inseparable (resp., orthogonal). Therefore, we will not distinguish  $p$ -equivalent matrices in this paper. In addition, for every statement in terms of the columns of a matrix  $A$ , there is a corresponding statement in terms of the rows of  $A$ . So we will only state our results in terms of columns in most cases.

Let  $f_0(m, n)$  (respectively,  $F_0(m, n)$ ) denote the smallest (respectively, largest) number of 1's in an  $m \times n$  inseparable orthogonal matrix of type 0 over  $\mathbb{Z}_2$ , and  $f_1(n)$  (resp.  $F_1(n)$ ) denote the smallest (resp., largest) number of 1's in an  $n \times n$  inseparable orthogonal matrix of type 1 over  $\mathbb{Z}_2$ . Clearly we have  $f_0(m, n) = f_0(n, m)$  and  $F_0(m, n) = F_0(n, m)$ . So we will assume  $m$  is odd if at least one of  $m$  and  $n$  is odd. The formulas for  $F_1(n)$  (Theorem 3.2),  $F_0(m, n)$  (Theorem 3.4),  $f_1(n)$  (Theorem 4.1), and  $f_0(m, n)$  (Theorem 4.2) are completely determined in this paper.

## 2. SOME PRELIMINARY RESULTS

We first note that the two matrices in Fig. 1 are both inseparable and of type 0. They will be used in constructing larger matrices in this paper.

LEMMA 2.1. *For any  $m \times n$  orthogonal matrix  $A$ , we have*

$$\omega(A) \equiv \begin{cases} 0 \pmod{4} & \text{if } A \text{ is of type 0} \\ n \pmod{4} & \text{if } A \text{ is a square matrix of type 1.} \end{cases}$$

*Proof.* Let  $k_i = \omega(C_i)$  for  $1 \leq i \leq n$ , and let  $\lambda_{ij}$  be the total number of columns covering both row  $i$  and row  $j$  for  $1 \leq i \neq j \leq m$ . Then by counting

$$A_{7,6} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \end{pmatrix} \quad A_7 = \begin{pmatrix} 1 & 1 & 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 & 1 & 0 \end{pmatrix}$$

(a)
(b)

FIGURE 1

the number  $s$  of submatrices of the form  $J_{2,1}$  (i.e., the number of columns in  $A$  covering an unordered pair of rows) in two different ways, it follows that

$$s = \sum_{i=1}^n \binom{k_i}{2} = \sum_{1 \leq i < j \leq m} \lambda_{ij}.$$

Since  $A$  is orthogonal, it follows that  $s$  is even. Therefore, we have

$$\sum_{i=1}^n (k_i^2 - k_i) = 2s \equiv 0 \pmod{4},$$

which implies that

$$\omega(A) = \sum_{i=1}^n k_i \equiv \sum_{i=1}^n k_i^2 \pmod{4}.$$

Note that either every  $k_i$  is even or every  $k_i$  is odd. Therefore, we have

$$\omega(A) \equiv \begin{cases} 0 \pmod{4} & \text{if } A \text{ is of type 0} \\ n \pmod{4} & \text{if } A \text{ is a square matrix of type 1.} \end{cases} \blacksquare$$

LEMMA 2.2. *For any  $m \times n$  orthogonal matrix  $A$  and any column  $C$  in  $A$ , we have*

$$\tau(C) \equiv \begin{cases} 0 \pmod{4} & \text{if } A \text{ is of type 0} \\ 1 \pmod{4} & \text{if } A \text{ is a square matrix of type 1.} \end{cases}$$

*Proof.* Let  $C$  be an arbitrary column in  $A$ . Define  $p_i = \omega(C, C_i)$  for  $1 \leq i \leq n$ . Without loss of generality, we assume  $C = C_1$  and  $\Omega(C_1) = \{R_i \mid 1 \leq i \leq p_1\}$ . Observe that  $p_i$  is even for each  $i$ ,  $2 \leq i \leq n$ , since  $A$  is orthogonal. We now define  $\lambda_{ij}$  to be the total number of columns in  $A$  covering both row  $i$  and row  $j$  for  $1 \leq i \neq j \leq p_1$ . Then by counting the number  $s$  of columns in  $A$  covering an unordered pair of rows in  $\Omega(C)$  in two different ways, it follows that

$$s = \sum_{i=1}^n \binom{p_i}{2} = \sum_{1 \leq i < j \leq p_1} \lambda_{ij}.$$

Since  $A$  is orthogonal, it follows that  $s$  is even. Therefore, we have

$$\sum_{i=1}^n (p_i^2 - p_i) = 2s \equiv 0 \pmod{4},$$

which implies that

$$\tau(C) = \sum_{i=1}^n p_i \equiv \sum_{i=1}^n p_i^2 \equiv p_1^2 \pmod{4}.$$

Therefore, we have

$$\tau(C) \equiv \begin{cases} 0 \pmod{4} & \text{if } A \text{ is of type 0} \\ 1 \pmod{4} & \text{if } A \text{ is a square matrix of type 1.} \end{cases} \blacksquare$$

The following result can be easily verified.

**LEMMA 2.3.** *Let  $A \neq I_1$  be an  $m \times n$  admissible orthogonal matrix such that  $m$  is odd. If  $A$  is either inseparable or of type 0, then we have  $m \geq 7$  and  $n \geq 6$ .*

For the remainder of this paper, we will assume that  $m \geq 7$  and  $n \geq 6$ , if  $m$  is odd.

**LEMMA 2.4.** *Let  $A$  be a  $7 \times n$  admissible orthogonal matrix of type 0, then the weight of every column of  $A$  is exactly 4.*

*Proof.* It is easy to see that  $A$  does not contain a column of weight 6 since  $A$  contains no zero row. Suppose  $A$  contains a column of weight 2, and let  $t$  be the maximum number of identical columns of weight 2. Then we have

$$A = \begin{pmatrix} J_{2,t} & B_1 \\ 0 & B_2 \end{pmatrix},$$

where  $B_2$  is a  $5 \times (n-t)$  orthogonal matrix of type 0. So  $B_2$  contains a 0 column by Lemma 2.3. But this is impossible since  $A$  is admissible and  $t$  is maximal. Therefore, the weight of every column of  $A$  is exactly 4.  $\blacksquare$

The following result is proved in [6].

**LEMMA 2.5.** *If there exists an inseparable orthogonal matrix  $A$  of odd order  $n$  such that the weight of every line in  $A$  is exactly 4, then  $n = 7$  and  $A$  is  $p$ -equivalent to  $A_7$ .*

Let  $A = (a_{ij})$  be a square matrix of order  $m$  such that  $a_{m1} = 1$ , and  $B = (b_{ij})$  be a square matrix of order  $n$  such that  $b_{1n} = 1$ . We define a square matrix  $C = (c_{ij}) = A \oplus B$  of order  $m+n-1$  as follows:

$$c_{ij} = \begin{cases} a_{i(j-n+1)} & \text{if } i \leq m \text{ and } j \geq n \\ b_{(i-m+1)j} & \text{if } i \geq m \text{ and } j \leq n \\ 0 & \text{if } i > m \text{ and } j > n \\ 1 & \text{if } i < m; j < n; \text{ and } a_{i1} = b_{1j} = 1 \\ 0 & \text{if } i < m; j < n; \text{ and either } a_{i1} = 0 \text{ or } b_{1j} = 0. \end{cases}$$

or equivalently, suppose

$$A = \begin{pmatrix} A_1 & A_2 \\ 1 & A_3 \end{pmatrix}, \quad B = \begin{pmatrix} B_1 & 1 \\ B_2 & B_3 \end{pmatrix}, \quad \text{and}$$

$Q = (q_{ij})$  is an  $(m-1) \times (n-1)$  matrix such that  $q_{ij} = 1$  if and only if  $a_{i1} = b_{1j} = 1$ . Then we define

$$A \oplus B = \begin{pmatrix} Q & A_1 & A_2 \\ B_1 & 1 & A_3 \\ B_2 & B_3 & O \end{pmatrix}.$$

LEMMA 2.6. Let  $A = (a_{ij})$  be an inseparable orthogonal matrix of order  $m$  and type 1 such that  $a_{m1} = 1$ , and  $B = (b_{ij})$  be an inseparable orthogonal matrix of order  $n$  and type 1 such that  $b_{1n} = 1$ . Then  $A \oplus B$  is an inseparable orthogonal matrix of order  $m+n-1$  and type 1. Moreover, we have  $\omega(A \oplus B) = \omega(A) + \omega(B) + ab - a - b$ , where  $a$  is the weight of the first column of  $A$  and  $b$  is the weight of the first row of  $B$ .

*Proof.* It can be easily seen that  $A \oplus B$  is an inseparable orthogonal matrix of order  $m+n-1$  and type 1. We also have

$$\begin{aligned} \omega(A \oplus B) &= \omega(A) + \omega(B) - 1 + (a-1)(b-1) \\ &= \omega(A) + \omega(B) + ab - a - b. \quad \blacksquare \end{aligned}$$

### 3. UPPER BOUNDS

#### 3.1. Matrices of Type 1

LEMMA 3.1. Let  $A$  be a matrix of order  $n$  and type 1, where  $n \geq 7$  is odd, such that the weight of every line of  $A$  is at least 3. Then we have  $\omega(A) \leq n^2 - 5n + 13$ .

*Proof.* We use induction on  $n$ . Clearly  $A$  contains no two identical lines, and the weight of each line of  $A$  is at most  $n-2$ . We may assume that  $n \geq q$ , since the case  $n=7$  can be easily verified. Now let  $r = r(A)$

(respectively,  $c = c(A)$ ) denote the number of rows (respectively, columns) in  $A$  of weight  $n - 2$ . We consider three cases.

*Case 1.*  $r(A) = c(A) = 0$ .

If the weight of every column is at most  $n - 6$ , then we have  $\omega(A) \leq n(n - 6) \leq n^2 - 5n + 13$ . So we assume  $A$  contains a column  $C_1$  with weight  $n - 4$ . Without loss of generality, we assume  $\Omega(C_1) = \{R_5, R_6, \dots, R_n\}$ . Then each remaining column of  $A$  covers at most three of the first 4 rows in  $A$ . So the number of 1's among the first 4 rows of  $A$  is at most  $3(n - 1)$ . Note that the weight of each row of  $A$  is at most  $n - 4$ . Therefore, we have

$$\omega(A) \leq 3(n - 1) + (n - 4)(n - 4) = n^2 - 5n + 13.$$

*Case 2.*  $r(A) \leq 3$ , and  $c(A) \leq 3$ .

Without loss of generality, we assume  $\Omega(C_1) = \{R_3, R_4, \dots, R_n\}$ . Then each remaining column of  $A$  covers exactly one of the first 2 rows in  $A$ . So the number of 1's among the first 2 rows of  $A$  is exactly  $n - 1$ . Note that the number of 1's among the remaining  $n - 2$  rows of  $A$  is at most  $(n - 4)(n - 2) + 6$  since  $r(A) \leq 3$ . Therefore, we have  $\omega(A) \leq n - 1 + (n - 4)(n - 2) + 6 = n^2 - 5n + 13$ .

*Case 3.*  $r(A) \geq 4$ , or  $c(A) \geq 4$ .

Clearly, we can assume that  $c(A) \geq 4$ , and  $A$  has the following format:

$$A = \left( \begin{array}{cc|cc} 1 & 0 & & \\ 0 & 1 & J_{2,t} & O \\ 0 & 0 & O & J_{1,(n-t-2)} \\ \hline 1 & 1 & & \\ \vdots & \vdots & & \\ 1 & 1 & X & \end{array} \right).$$

Since  $c(A) \geq 4$ , it follows that if  $C_i$  is a column with weight  $n - 2$ , then  $i \leq t + 2$ . Without loss of generality, we assume that each of the first 4 columns of  $A$  has weight  $n - 2$ . Let  $B$  be the submatrix of order  $n - 2$  obtained from  $A$  by deleting the first two columns and first two rows from  $A$ . Then  $B$  is of type 1. If  $B$  contains a line of weight 1, then we can easily see that the columns  $C_3$  and  $C_4$  (of weight  $n - 2$  each) must be identical, which is impossible. Note that the third row of  $A$  has weight at least 3, which implies that  $t \leq n - 5$ . Therefore, by induction we have

$$\begin{aligned} \omega(A) &= \omega(B) + 2(n - 2) + 2t \\ &\leq (n - 2)^2 - 5(n - 2) + 13 + 2(n - 2) + 2(n - 5) \\ &= n^2 - 5n + 13. \quad \blacksquare \end{aligned}$$



THEOREM 3.2. *We have*

$$F_1(n) = \begin{cases} n^2 - n & \text{if } n \geq 4 \text{ is even} \\ n^2 - 5n + 13 & \text{if } n \geq 7 \text{ is odd.} \end{cases}$$

*Proof.* If  $A$  is an admissible matrix of order  $n$  and type 1, where  $n$  is even, then we have  $n \geq 4$  and the weight of each line of  $A$  is at most  $n - 1$ . So we have  $F_1(n) \leq n^2 - n$  if  $n \geq 4$  is even. It is easy to see that  $J_n - I_n$  is an inseparable orthogonal matrix of type 1 when  $n \geq 4$  is even. It then follows by Lemma 2.6 that  $(J_4 - I_4) \oplus (J_{n-3} - I_{n-3})$  is an inseparable orthogonal matrix of type 1 when  $n \geq 7$  is odd. ■

### 3.2. Matrices of Type 0

The complement of a matrix  $A$  is obtained from  $A$  by replacing every 0 by a 1 and every 1 by a 0.

LEMMA 3.3. *Let  $B$  be an  $m \times n$  matrix such that  $BB^T = J_m$  and  $B^TB = J_n$ , where  $m$  and  $n$  are both odd. If the number of 0's in every line of  $B$  is at least 2, then we have  $m \geq 7$ ,  $n \geq 7$ , and  $\omega(B) \geq 3(m + n - 7)$ .*

*Proof.* We use contradiction. Suppose there exists an  $n \times n$  matrix  $B$  such that  $\omega(B) < 3(m + n - 7)$ ,  $BB^T = J_m$  and  $B^TB = J_n$ , where  $m$  and  $n$  are both odd, and the number of 0's in every line of  $B$  is at least 2. We further assume that  $m + n$  is minimal among such matrices. Let  $A$  be the complement of  $B$ . Then  $A$  is an admissible matrix of type 0. So  $m \geq 7$  and  $n \geq 7$  by Lemma 2.3. Clearly the weight of each line of  $B$  is odd and at least 3. Let  $t_r = t_r(B)$  (respectively,  $t_c = t_c(B)$ ) denote the number of rows (respectively, columns) of weight 3 in  $B$ .

CLAIM 1.  *$B$  contains at most two rows of weight at least  $n - 4$  each, and  $B$  contains at most two columns of weight at least  $m - 4$  each.*

*Proof.* If  $B$  contains at least three rows of weight at least  $n - 4$  each, then we would have

$$\omega(B) \geq 3(n - 4) + (m - 3)3 = 3(m + n - 7).$$

which is impossible.

CLAIM 2. *If  $B$  contains two identical columns of weight 3 each, then  $t_c \leq 4$ .*

*Proof.* Let  $B_1$  be the submatrix of  $B$  obtained by deleting two identical columns of weight 3. Then  $B_1$  is an  $m \times (n - 2)$  matrix such that  $B_1 B_1^T = J_m$  and  $B_1^T B_1 = J_{n-2}$ . If the number of 0's in every line of  $B_1$  is at least 2, then

we would have  $\omega(B) = \omega(B_1) + 6 \geq 3(m+n-2-7) + 6 = 3(m+n-7)$ , which is impossible. So there exists a row in  $B_1$  of weight  $n-2$ , which means that  $B$  contains a row of weight  $n-2$ , and  $B$  contains no three identical columns of weight three.

Suppose  $B$  contains another pair of identical columns of weight three. Then  $B$  has the following format (by interchanging some lines in  $B$ ):

$$B = \left( \begin{array}{cccc|c} 0 & 0 & 1 & 1 & J_{2, (n-4)} \\ 1 & 1 & 0 & 0 & \\ \hline 1 & 1 & 1 & 1 & \\ 1 & 1 & 0 & 0 & B_2 \\ 0 & 0 & 1 & 1 & \\ \hline O & & & & \end{array} \right),$$

where  $B_2$  is an  $(m-2) \times (n-4)$  matrix such that  $B_2 B_2^T = J_{m-2}$  and  $B_2^T B_2 = J_{n-4}$ . If  $t_c \geq 5$ , then  $B_2$  contains a column of weight 1, which implies that the corresponding row of  $B_2$  has weight at least  $n-4$ . Therefore,  $B$  contains at least three rows of weight at least  $n-4$  each, which contradicts Claim 1. Therefore, we have  $t_c \leq 4$  if  $B$  contains two different pairs of identical columns of weight three.

Now assume  $B$  contains a unique pair of identical columns of weight three. Suppose  $t_c \geq 5$ , then  $B$  must have the following format:

$$B = \left( \begin{array}{ccccc|c} 0 & 0 & 1 & 1 & 1 & 1 \dots 1 \\ \hline & J_{3,2} & I_3 & & & X \\ & O & I_3 & & & \\ \hline & O & & & & Y \end{array} \right).$$

This implies that each column of  $B$  covers at least three of the first 7 rows of  $B$ . So the total number of 1's in the first 7 rows of  $B$  is at least  $3n$ . Therefore,

$$\omega(B) \geq 3n + 3(m-7) = 3(m+n-7),$$

which is impossible. This proves Claim 2.

**CLAIM 3.** *If  $t_c \geq 5$ , then  $B$  contains no two identical columns of weight 3, and there exists a row in  $B$  which covers every column of weight three in  $B$ .*

*Proof.* Assume  $t_c \geq 5$ . Then Claim 2 implies that  $B$  contains no two identical columns of weight 3. If  $B$  contains 4 columns of weight 3, each of

which covers a given row  $R$ , then every other column of  $B$  also covers row  $R$  by Claim 2, and we are done. So we assume that every row in  $B$  covers at most 3 columns of weight 3 in  $B$ . Then we can easily see that  $B$  has the following format:

$$B = \left( \begin{array}{cccccc|c} 1 & 1 & 1 & 0 & 0 & 0 & X \\ 1 & 0 & 0 & 1 & 1 & 0 & X \\ 1 & 0 & 0 & 0 & 0 & 0 & X \\ 0 & 1 & 0 & 1 & 0 & 0 & X \\ 0 & 1 & 0 & 0 & 1 & 0 & X \\ 0 & 0 & 1 & 1 & 0 & 0 & X \\ 0 & 0 & 1 & 0 & 1 & 0 & X \\ \hline & & & & & 0 & Y \end{array} \right).$$

Therefore, each column of  $B$  covers at least three of the first 7 rows of  $B$ . So the total number of 1's in the first 7 rows of  $B$  is at least  $3n$ . Therefore,

$$\omega(B) \geq 3n + 3(m-7) = 3(m+n-7),$$

which is impossible. This proves Claim 3.

Let  $r = r(B)$  (respectively,  $c = c(B)$ ) denote the minimum weight of a row (respectively, column) in  $B$ . Then it follows by Claim 1 that  $r \leq n-6$  and  $c \leq m-6$ .

CLAIM 4. *We have*

$$\omega(B) \geq rm + n - rc + c - 1 \quad \text{and} \quad \omega(B) \geq cn + m - rc + r - 1. \quad (1)$$

*Proof.* Without loss of generality, we assume that the weight of the first column  $C_1$  of  $B$  is minimal, and  $C_1$  consists of the first  $c$  rows of  $B$ . Then each of the remaining columns of  $B$  covers at least one of the first  $c$  rows of  $B$ . So the total number of 1's in the first  $c$  rows of  $B$  is at least  $c + (n-1)$ . Note that the weight of each row in  $B$  is at least  $r$ . Therefore,

$$\omega(B) \geq (c + n - 1) + r(m - c) = rm + n - rc + c - 1.$$

By symmetry, we have  $\omega(B) \geq cn + m - rc + r - 1$ .

CLAIM 5. *We have*

$$\omega(B) \geq 5m + n - 4c - 9 \quad \text{and} \quad \omega(B) \geq 5n + m - 4r - 9. \quad (2)$$

*Proof.* We will prove the first inequality only. Without loss of generality, we assume that the weight of the first column  $C_1$  of  $B$  is minimal, and  $\Omega(C_1)$  consists of the first  $c$  rows of  $B$ . Then each of the remaining columns of  $B$  covers at least one of the first  $c$  rows of  $B$ . So the total number of 1's in the first  $c$  rows of  $B$  is at least  $c + (n - 1)$ . Let  $\mathfrak{R}$  be the set of remaining  $m - c$  rows of  $B$ . If there are at most 4 rows of weight 3 in  $\mathfrak{R}$ , then we have

$$\omega(B) \geq n + c - 1 + 5(m - c) - 8 = 5m + n - 4c - 9.$$

So we assume that there are at least 5 rows of weight 3 in  $\mathfrak{R}$ . Let  $X_1$  be the set of rows of weight 3 in  $\mathfrak{R}$ . Then we may assume that each row of weight 3 covers the last column  $C_n$  of  $B$  by Claim 3. Let  $X_2$  be the set of rows of weight at least 5 in  $\mathfrak{R}$  covering the last column of  $B$ , and  $X_3 = \mathfrak{R} \setminus (X_1 \cup X_2)$ . Also let  $x_i = |X_i|$  for  $1 \leq i \leq 3$ . Note that  $x_1 \geq 5$ ,  $x_1 + x_2 + x_3 = m - c$ , and the weight of each row in  $X_3$  is at least  $x_1$ . Therefore, we have

$$\begin{aligned} \omega(B) &\geq n + c - 1 + 3x_1 + 5x_2 + x_1x_3 \\ &= n + c - 1 + 5(m - c) + (x_1 - 5)x_3 - 2x_1. \end{aligned}$$

If  $x_3 \geq 2$ , then we have

$$\omega(B) \geq n + c - 1 + 5(m - c) - 10 = 5m + n - 4c - 11.$$

Since  $\omega(B) \equiv mn \pmod{4}$  by Lemma 2.1, it follows that  $\omega(B) \geq 5m + n - 4c - 9$  if  $x_3 \geq 2$ . Now assume  $x_3 \leq 1$ . We also assume  $R_n \notin \Omega(C_n)$  if  $x_3 = 1$ . Then we have  $x_2 \geq m - c - x_1 - 1$ . Recall that  $\Omega(C_1) = \{R_1, R_2, \dots, R_c\}$  is the set of the first  $c$  rows of  $B$ . Now let  $Y = \Omega(C_n) \cap \Omega(C_1) = \{R_1, R_2, \dots, R_y\}$ , where  $y = |Y|$ ,  $Z = \Omega(C_1) \setminus Y$ , and  $z = |Z|$ . Then we have  $y = c - z \leq c - 2$  since the number of 0's in the last column of  $B$  is at least two and  $z$  is even. Given any column  $C$  in  $B$ , we have  $\omega(C, C_1)$  and  $\omega(C, C_n)$  are both odd, which implies that  $|\Omega(C) \cap Y|$  is odd if  $x_3 = 0$  and  $|\Omega(C) \cap (Y \cup R_n)|$  is odd if  $x_3 = 1$ . So the total number of 1's among the first  $y$  rows (and the last row  $R_n$  if  $x_3 = 1$ ) in  $B$  is at least  $2y + (n - 2)$ . Note that the weight of each row in  $Z$  is at least  $(x_1 + 1)$ . Therefore, we have

$$\begin{aligned} \omega(B) &\geq 2y + (n - 2) + (x_1 + 1)z + 3x_1 + 5x_2 \\ &\geq 2y + (n - 2) + (x_1 + 1)(c - y) + 3x_1 + 5(m - c - x_1 - 1) \\ &= 5m + n - 4c + y - 7 + (c - 2 - y)x_1 \\ &\geq 5m + n - 4c + y - 7 \\ &> 5m + n - 4c - 9. \end{aligned}$$

Similarly, we can prove that  $\omega(B) \geq 5n + m - 4r - 9$ . This proves Claim 5.

We now consider 3 cases.

*Case 1.*  $r \geq 5$  and  $c \geq 5$ .

Since  $r \leq n-6$  and  $c \leq m-6$ , it follows by Claim 4 that

$$\begin{aligned} 2\omega(B) &\geq (r+1)m + (c+1)n - 2rc + r + c - 2 \\ &= 6(m+n) - 42 + (r-5)m + (c-5)n - 2rc + r + c + 40 \\ &= 6(m+n) - 42 + (r-5)(m-c-4) + (c-5)(n-r-4) \\ &\geq 6(m+n) - 42. \end{aligned}$$

Therefore, we have  $\omega(B) \geq 3(m+n-7)$ , which is a contradiction.

*Case 2.*  $r = 3$  and  $c \geq 5$ .

It follows by Claims 4 and 5 that

$$\begin{aligned} 2\omega(B) &\geq (cn + m - 3c + 2) + (5m + n - 4c - 9) \\ &= 6m + (c+1)n - 7c - 7 \\ &= 6m + 6n - 42 + (c-5)(n-7) \\ &\geq 6(m+n-7). \end{aligned}$$

Therefore, we have  $\omega(B) \geq 3(m+n-7)$ , which is a contradiction.

*Case 3.*  $r = c = 3$ .

it follows by Claim 5 that

$$2\omega(B) \geq (5m + n - 21) + (5n + m - 21) = 6(m+n-7).$$

Therefore, we have  $\omega(B) \geq 3(m+n-7)$ , which is a contradiction. ■

**THEOREM 3.4.** *We have*

$$F_0(m, n) = \begin{cases} mn & \text{if } m \text{ and } n \text{ are both even} \\ (m-3)n & \text{if } m \geq 7 \text{ is odd and } n \geq 6 \text{ is even} \\ mn - 3m - 3n + 21 & \text{if } m \geq 7 \text{ and } n \geq 7 \text{ are both odd.} \end{cases}$$

*Proof.* Clearly we have  $F_0(m, n) = mn$  if  $m$  and  $n$  are both even. Let  $A$  be an  $m \times n$  admissible orthogonal matrix of type 0, where  $m$  is odd. Then the weight of each column of  $A$  is at most  $m-3$ . So  $\omega(A) \leq (m-3)n$ . This

implies that  $F_0(m, n) \leq (m-3)n$  if  $m$  is odd. So it suffices to construct the upper bound by Lemma 3.3. We define

$$A = \left( \begin{array}{c|c} A_{7,6} & \begin{matrix} J_{4,(n-6)} \\ O \end{matrix} \\ \hline J_{(m-7),6} & J_{(m-7),(n-6)} \end{array} \right), \quad \text{where}$$

$A_{7,6}$  is as in Fig. 1a. Then  $A$  is an  $m \times n$  inseparable orthogonal matrix of type 0 with  $\omega(A) = (m-3)n$ , if  $m \geq 7$  is odd and  $n \geq 6$  is even.

We now define

$$B = \left( \begin{array}{c|c} A_7 & \begin{matrix} J_{4,(n-7)} \\ O \end{matrix} \\ \hline J_{(m-7),4} & J_{(m-7),(n-7)} \end{array} \right), \quad \text{where}$$

$A_7$  is as in Fig. 1b. Then  $B$  is an  $m \times n$  inseparable orthogonal matrix of type 0 with  $\omega(A) = mn - 3m - 3n + 21$ , if  $m \geq 7$  and  $n \geq 7$  are both odd. ■

## 4. LOWER BOUNDS

### 4.1. Matrices of Type 1

**THEOREM 4.1.** *We have  $f_1(n) = 5n - 8$  if  $n \geq 4$  and  $n \neq 5$ .*

*Proof.* It is easy to see that  $(\begin{smallmatrix} I_{n-2} & J_{I_2}^{(n-2),2} \\ J_{2,(n-2)} & I_2 \end{smallmatrix})$  is an inseparable orthogonal matrix of type 1 when  $n \geq 4$  is even, and  $(J_4 - I_4) \oplus (\begin{smallmatrix} I_{n-5} & J_{I_2}^{(n-5),2} \\ J_{2,n-5} & I_2 \end{smallmatrix})$  is an inseparable orthogonal matrix of type 1 when  $n \geq 7$  is odd.

We now prove the lower bound. It is easy to verify the result for  $n \leq 5$ . So we assume  $n \geq 6$ . Suppose  $n \geq 6$  is the smallest integer such that  $f_1(n) < 5n - 8$ , and let  $A$  be an inseparable orthogonal matrix of order  $n$  and type 1 such that  $\omega(A) = f_1(n)$ .

**CLAIM 1.**  *$A$  does not contain three columns  $C_1, C_2$ , and  $C_3$  of weight 3 each such that  $\omega(C_1, C_2) = \omega(C_1, C_3) = \omega(C_2, C_3) = 2$ . Moreover, if  $A$  contains two columns  $C_1$  and  $C_2$  of weight 3 each such that  $\omega(C_1, C_2) = 2$ , then no column (other than  $C_1$  and  $C_2$ ) in  $A$  covers both rows in  $\Omega(C_1) \cap \Omega(C_2)$ .*

*Proof.* First assume that  $A$  contains two columns  $C_1$  and  $C_2$  of weight 3 each such that  $\omega(C_1, C_2) = 2$ ,  $\Omega(C_1) = \{R_1, R_3, R_4\}$ , and  $\Omega(C_2) = \{R_2, R_3, R_4\}$ . Suppose there exists another column  $C$  covering both  $R_3$  and  $R_4$ . Then we can assume that  $A$  has the following format:

$$A = \left( \begin{array}{cc|cccc} 1 & 0 & 1 & \dots & 1 & 0 & \dots & 0 \\ 0 & 1 & 1 & \dots & 1 & 0 & \dots & 0 \\ \hline 1 & 1 & & & & & & \\ 1 & 1 & & & & & & \\ 0 & 0 & & & B & & & \\ \vdots & \vdots & & & & & & \\ 0 & 0 & & & & & & \end{array} \right).$$

Then it can be seen that  $B$  is an inseparable orthogonal matrix of order  $n-2$  and type 1. Therefore, we have

$$\omega(A) \geq \omega(B) + 10 \geq 5(n-2) - 8 + 10 = 5n - 8,$$

which is impossible. Therefore,  $C_1$  and  $C_2$  are the only columns in  $A$  covering both  $R_3$  and  $R_4$ .

Now suppose  $C_3$  is another column in  $A$  with weight 3 such that  $\omega(C_1, C_2) = \omega(C_1, C_3) = \omega(C_2, C_3) = 2$ . Then  $A$  has the following format:

$$A = \left( \begin{array}{ccc|cc} 1 & 1 & 1 & & \\ 1 & 1 & 0 & O & O \\ 1 & 0 & 1 & J_{3,t} & O \\ 0 & 1 & 1 & & \\ \hline & & & O & X \end{array} \right),$$

where  $t \geq 3$  is odd. Let  $B$  be the submatrix obtained by deleting the first 3 columns and the first 3 rows from  $A$ . Then it can be seen that  $B$  is an inseparable orthogonal matrix of order  $n-3$  and type 1. Therefore, we have

$$\omega(A) = \omega(B) + 2t + 9 \geq 5(n-3) - 8 + 15 = 5n - 8,$$

which is impossible. Therefore,  $A$  does not contain three columns  $C_1$ ,  $C_2$ , and  $C_3$  with weight 3 each such that  $\omega(C_1, C_2) = \omega(C_1, C_3) = \omega(C_2, C_3) = 2$ .

**CLAIM 2.** Assume  $A$  contains two columns  $C_1$  and  $C_2$  of weight 3 each such that  $\omega(C_1, C_2) = 2$ . If  $\Omega(C_1) = \{R_1, R_3, R_4\}$  and  $\Omega(C_2) = \{R_2, R_3, R_4\}$ , then we have  $\omega(R_1) = \omega(R_2) \geq 7$ ,  $\omega(R_3) \geq 5$ , and  $\omega(R_4) \geq 5$ .

*Proof.* Assume that  $A$  contains two columns  $C_1$  and  $C_2$  of weight 3 each such that  $\omega(C_1, C_2) = 2$ ,  $\Omega(C_1) = \{R_1, R_3, R_4\}$ , and  $\Omega(C_2) = \{R_2, R_3, R_4\}$ . Then we have

$$A = \left( \begin{array}{cc|cc} 1 & 0 & 1 \dots 1 & 0 \dots 0 \\ 0 & 1 & 1 \dots 1 & 0 \dots 0 \\ \hline 1 & 1 & & \\ 1 & 1 & & \\ 0 & 0 & & B \\ \vdots & \vdots & & \\ 0 & 0 & & \end{array} \right),$$

where  $B$  is separable according to the proof of Claim 1. Therefore,  $A$  has the following format:

$$A = \left( \begin{array}{cc|cc|cc} 1 & 0 & & & & & \\ 0 & 1 & J_{2,\alpha} & O & J_{2,\beta} & O & \\ \hline 1 & 1 & & & & & \\ 0 & 0 & J_{1,\alpha} & O & & O & \\ \vdots & \vdots & X & & & & \\ 0 & 0 & & & & & \\ \hline 1 & 1 & & & & & \\ 0 & 0 & O & & J_{1,\beta} & O & \\ \vdots & \vdots & & & Y & & \\ 0 & 0 & & & & & \end{array} \right),$$

where both  $\alpha$  and  $\beta$  are odd. If  $\alpha = 1$ , then it can be seen that the weight of the third column of  $A$  is 3, which is impossible by Claim 1. So we have  $\alpha \geq 3$ . Similarly, we have  $\beta \geq 3$ . This implies that  $\omega(R_1) = \omega(R_2) = \alpha + \beta + 1 \geq 7$ ,  $\omega(R_3) = \alpha + 2 \geq 5$ , and  $\omega(R_4) = \beta + 2 \geq 5$ .

**CLAIM 3.** *If  $A$  contains a column  $C_1$  of weight 3 and a row  $R_1$  of weight 3 such that  $\Omega(C_1) = \{R_1, R_2, R_3\}$  and  $\Omega(R_1) = \{C_1, C_2, C_3\}$ , then we have  $\omega(R_2) = \omega(R_3) \geq 5$ ,  $\omega(C_2) = \omega(C_3) \geq 5$ , and  $\omega(C_2) + \omega(R_2) \geq 12$ .*

*Proof.* Assume that  $A$  contains a column  $C_1$  of weight 3 and a row  $R_1$  of weight 3 such that  $\Omega(C_1) = \{R_1, R_2, R_3\}$  and  $\Omega(R_1) = \{C_1, C_2, C_3\}$ . Then  $A$  has the following format:



$$A = \left( \begin{array}{ccc|ccc} 1 & 1 & 1 & 0 \dots 0 & 0 \dots 0 & \\ 1 & 1 & 0 & 1 \dots 1 & 0 \dots 0 & \\ 1 & 0 & 1 & 1 \dots 1 & 0 \dots 0 & \\ \hline 0 & 1 & 1 & & & \\ \vdots & \vdots & & & & \\ 0 & 1 & 1 & & & \\ 0 & 0 & 0 & & & \\ \vdots & \vdots & & & & \\ 0 & 0 & 0 & & & \end{array} \right),$$

$B$

where  $\omega(R_2) = \omega(R_3)$  and  $\omega(C_2) = \omega(C_3)$ . It then follows by Claim 1 that  $\omega(R_2) = \omega(R_3) \geq 5$  and  $\omega(C_2) = \omega(C_3) \geq 5$ . Clearly,  $B$  is an orthogonal matrix of order  $n - 3$  and type 1. If  $B$  were inseparable, then we would have

$$\omega(A) \geq \omega(B) + 19 \geq 5(n - 3) - 8 + 19 > 5n - 8,$$

which is impossible. So  $B$  is separable.

Now suppose  $\omega(R_2) = \omega(R_3) = 5$  and  $\omega(C_2) = \omega(C_3) = 5$ . Then  $A$  has the following format:

$$A = \left( \begin{array}{ccc|ccc|ccc} 1 & 1 & 1 & 0 & 0 \dots 0 & 0 & 0 & 0 \dots 0 & \\ 1 & 1 & 0 & 1 & 0 \dots 0 & 1 & 1 & 0 \dots 0 & \\ 1 & 0 & 1 & 1 & 0 \dots 0 & 1 & 1 & 0 \dots 0 & \\ \hline 0 & 1 & 1 & & & & & & \\ 0 & 0 & 0 & & & & & & \\ \vdots & \vdots & \vdots & & & & & & \\ 0 & 0 & 0 & & & & & & \\ \hline 0 & 1 & 1 & & & & & & \\ 0 & 1 & 1 & & & & & & \\ 0 & 0 & 0 & & & & & & \\ \vdots & \vdots & \vdots & & & & & & \\ 0 & 0 & 0 & & & & & & \end{array} \right).$$

$A_1$        $O$

$O$        $A_2$

This would imply that both the fourth column and the fourth row of  $A$  have weight 3, which is impossible by Claim 1. Therefore, we have  $\omega(C_2) + \omega(R_2) \geq 12$ .

CLAIM 4. *If  $A$  contains a column  $C$  of weight 3 such that each row in  $\Omega(C)$  has weight at least 5, then we have  $\tau(C) \geq 17$ .*

*Proof.* Assume  $\Omega(C) = \{R_1, R_2, R_3\}$ . Then we have  $\tau(C) = \omega(R_1) + \omega(R_2) + \omega(R_3) \geq 15$ . Note that  $\tau(C) \equiv 1 \pmod{4}$  by Lemma 2.2. Therefore, we have  $\tau(C) \geq 17$ .

Now let  $\mathfrak{R}$  (respectively,  $\mathfrak{C}$ ) be the set of rows (respectively, columns) of weight 3 in  $A$ . A row  $R$  (respectively, a column  $C$ ) in  $\mathfrak{R}$  (respectively,  $\mathfrak{C}$ ) is said to be *good* if there exists another row  $R^*$  (respectively, column  $C^*$ ) in  $\mathfrak{R}$  (respectively,  $\mathfrak{C}$ ) such that  $\omega(R, R^*) > 0$  (respectively,  $\omega(C, C^*) > 0$ ). We then let  $\mathfrak{R}_1$  (respectively,  $\mathfrak{C}_1$ ) be the set of good rows (respectively, good columns) in  $\mathfrak{R}$  (respectively,  $\mathfrak{C}$ );  $\mathfrak{R}_2$  (respectively,  $\mathfrak{C}_2$ ) be the set of rows (respectively, columns) in  $\mathfrak{R} \setminus \mathfrak{R}_1$  (respectively,  $\mathfrak{C} \setminus \mathfrak{C}_1$ ) such that each of the corresponding three columns (respectively, rows) has weight at least 5;  $\mathfrak{R}_3$  be the set of rows in  $\mathfrak{R} \setminus (\mathfrak{R}_1 \cup \mathfrak{R}_2)$  such that the total number of 1's in the corresponding three columns is at least 17; and  $\mathfrak{C}_3$  be the set of columns in  $\mathfrak{C} \setminus (\mathfrak{C}_1 \cup \mathfrak{C}_2)$  that does not cover any row in  $\mathfrak{R}_3$ . We also define  $r_i = |\mathfrak{R}_i|$  and  $c_i = |\mathfrak{C}_i|$  for  $1 \leq i \leq 3$ . Then by counting the number of 1's in each column of  $A$ , we have

$$\omega(A) \geq 5n - 2(c_1 + c_2 + c_3) + 2r_1 + 2r_2 + 2r_3. \quad (3)$$

Similarly, by counting the number of 1's in each row of  $A$ , we have

$$\omega(A) \geq 5n - 2(r_1 + r_2 + r_3) + 2c_1 + 2c_2 + 2c_3. \quad (4)$$

By adding (3) and (4) together, it follows that  $\omega(A) \geq 5n$ , which is a final contradiction. ■

## 4.2. Matrices of Type 0

THEOREM 4.2. *We have*

$$f_0(m, n) = \begin{cases} 2(m+n) - 4 & \text{if } m \text{ and } n \text{ are both even} \\ \text{does not exist} & \text{if either } m \leq 5 \text{ or } n \leq 5; \text{ and } m \text{ is odd} \\ 4n & \text{if } m = 7 \text{ and } n \geq 6 \\ 2(m+n) - 2 & \text{if } m \geq 9 \text{ is odd and } n \geq 6 \text{ is even} \\ 2(m+n) + 4 & \text{if } m \text{ and } n \text{ are both odd; } m \leq n; \text{ and } m = 9 \text{ or } 11 \\ 2(m+n) & \text{if } m \geq 13; n \geq 13; \text{ and } m \text{ and } n \text{ are both odd.} \end{cases}$$

*Proof.* We first prove the upper bound. Recall that  $A_{7,6}$  is the  $7 \times 6$  matrix in Fig. 1a and  $A_7$  is the  $7 \times 7$  matrix in Fig. 1b. If  $m$  and  $n$  are both even, then we let  $A$  be the matrix in Fig. 2a. If  $m \geq 9$  is odd and  $n \geq 6$  is

$$\begin{array}{ccc}
 \begin{pmatrix} J_{2,(n-2)} & J_2 \\ O & J_{(m-2),2} \end{pmatrix} & & \left( \begin{array}{cc|cc} J_{2,(n-8)} & J_2 & J_2 & O \\ O & J_{(m-9),2} & O & O \\ \hline O & & A_{7,6} & \end{array} \right) \\
 \text{(a)} & & \text{(b)}
 \end{array}$$

FIGURE 2

even, then we let  $A$  be the matrix in Fig. 2b. If  $m = 7$  and  $n \geq 6$  is even, then we let  $A$  be the matrix in Fig. 3a. If  $m = 7$  and  $n \geq 7$  is odd, then we let  $A$  be the matrix in Fig. 3b.

We now assume  $m$  and  $n$  are both odd and  $9 \leq m \leq n$ . It can be easily seen that the following matrix  $A_9$  is an inseparable orthogonal matrix of type 0 and order 9.

$$A_9 = \left( \begin{array}{ccc|ccc|ccc} 0 & 0 & 1 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 1 & 1 & 0 & 1 & 1 & 0 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 & 1 & 1 & 0 & 1 & 1 \\ \hline 0 & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 1 \end{array} \right).$$

If  $m = 9$ , then we let  $A$  be the matrix in Fig. 4a. If  $m = 11$ , then we let  $A$  be the matrix in Fig. 4b. If  $m \geq 13$ , then we let  $A$  be the matrix in Fig. 5. Then we can easily verify that the matrix  $A$  constructed above is an  $m \times n$  inseparable orthogonal matrix of type 0 that achieves the bound given in Theorem 4.2.

We now prove the lower bound. Lemma 1.3 implies that  $f_0(m, n) \geq 2(m + n) - 4$ . In particular, we have  $f_0(m, n) \geq 2(m + n) - 2$  if  $m$  is odd and  $n$  is even by Lemma 2.1. Lemma 2.4 implies that  $f_0(m, n) \geq 4n$  if  $m = 7$  and  $n \geq 6$ . We now assume  $m \geq 9$  and  $n \geq 9$  are both odd. Suppose

$$\begin{array}{ccc}
 \left( A_{7,6} \mid \begin{array}{c} J_{4,(n-6)} \\ O \end{array} \right) & & \left( A_7 \mid \begin{array}{c} J_{4,(n-7)} \\ O \end{array} \right) \\
 \text{(a)} & & \text{(b)}
 \end{array}$$

FIGURE 3

$$\begin{array}{ccc}
 \left( A_9 \mid \begin{array}{c} J_{2,(n-9)} \\ O \end{array} \right) & & \left( \begin{array}{c|cc} A_9 & J_2 & O \\ \hline & O & O \\ \hline O & J_{2,(n-9)} & \end{array} \right) \\
 \text{(a)} & & \text{(b)}
 \end{array}$$

FIGURE 4

Theorem 4.2 is false. Then it follows by Lemma 2.1 that there exists an  $m \times n$  inseparable matrix  $A$  of type 0 such that (i)  $\omega(A) \leq 2(m+n)$ ; (ii)  $\omega(A) < 2(m+n)$  if  $m \geq 13$  and  $n \geq 13$ ; and (iii)  $m+n$  is minimal among such matrices.

**CLAIM 1.**  *$A$  contains a line of weight 2. In particular, if  $m < n$ , then  $A$  contains a column of weight 2.*

*Proof.* Suppose not. Then we have  $\omega(A) \geq 4n > 2(m+n)$  if  $n > m$ , which contradicts the choice of  $A$ . So we assume  $n = m$ . Then at least one line of  $A$  has weight larger than 4 by Lemma 2.5, which implies that  $\omega(A) > 4n = 2(m+n)$ , which is impossible.

**CLAIM 2.**  *$A$  contains no two identical lines of weight 2.*

*Proof.* Suppose not, and let  $B$  be a submatrix of  $A$  by deleting two identical columns of weight 2. Then  $B$  is an  $m \times (n-2)$  inseparable matrix of type 0. Since  $m+n$  is minimal, it follows that  $n-2 = 7$ , which implies that  $m \geq n = 9$ . It then follows by Lemma 2.4 that

$$\omega(A) = \omega(B) + 4 \geq 4m + 4 > 2(m+n).$$

This proves Claim 2.

We now consider 2 cases.

*Case 1.*  $n \geq m$  and  $m = 9$  or  $11$ .

In this case, we can assume that  $A$  contains a column  $C_1$  of weight 2 by Claim 1. Note that  $\omega(A) \leq 2(m+n) \leq 4n$ . Now let  $B$  be the submatrix

$$\left( \begin{array}{cc|cc|cc} J_{2,(n-13)} & J_2 & J_2 & O & & \\ O & J_{(m-13),2} & O & O & & \\ \hline & O & J_2 & O & A_{7,6}^T & \\ & & O & O & \hline & & A_{7,6} & O & & \end{array} \right)$$

FIGURE 5

obtained from  $A$  by deleting column  $C_1$  and the corresponding two rows. If  $m=9$ , then  $B$  is an admissible  $7 \times (n-1)$  orthogonal matrix of type 0 by Claim 2. It then follows by Lemma 2.4 that

$$\omega(A) \geq \omega(B) + 8 \geq 4(n-1) + 8 = 4n + 4,$$

which is impossible. So we assume  $m=11$ .

If  $A$  contains 2 columns of weight 2, then  $A$  has the following format:

$$A = \left( \begin{array}{cc|cc|c} 1 & 0 & & & \\ 1 & 0 & & & \\ 0 & 1 & J_{4,t} & & B_1 \\ 0 & 1 & & & \\ \hline & & O & O & B_2 \end{array} \right),$$

where  $B_2$  has no zero column. Therefore,  $B_2$  is an admissible  $7 \times (n-t-2)$  orthogonal matrix of type 0. It can be easily seen that  $\omega(B_1) \geq 6$ . It then follows by Lemma 2.4 that

$$\omega(A) \geq \omega(B_1) + \omega(B_2) + 4t + 4 \geq 6 + 4(n-t-2) + 4t + 4 = 4n + 2,$$

which is a contradiction. So  $A$  contains exactly one column of weight 2.

If every column  $C$  (other than  $C_1$ ) in  $A$  such that  $\Omega(C_1) \subseteq \Omega(C)$  has weight at least 6, then we would have  $\omega(A) \geq 4n - 2 + 6 = 4n + 4$ , which is impossible. So there exists a column, say  $C_2$ , of weight 4 in  $A$  such that  $\Omega(C_1) \subseteq \Omega(C_2)$ , and  $A$  has the following format:

$$A = \left( \begin{array}{cc|cc|c} 1 & 1 & & & \\ 1 & 1 & & & \\ 0 & 1 & J_{4,t} & & B_1 \\ 0 & 1 & & & \\ \hline & & O & O & B_2 \end{array} \right),$$

where  $B_2$  has no zero column. Therefore,  $B_2$  is an admissible  $7 \times (n-t-2)$  orthogonal matrix of type 0. It can be easily seen that  $\omega(B_1) \geq 4$ . It then follows by Lemma 2.4 that

$$\omega(A) \geq \omega(B_1) + \omega(B_2) + 4t + 6 \geq 4 + 4(n-t-2) + 4t + 6 = 4n + 2,$$

which is again a contradiction.

*Case 2.*  $m \geq 13$ , and  $n \geq 13$ .

We first assume that  $A$  contains a column  $C_1$  of weight 2 such that  $\tau(C_1) = 8$ . Let  $B$  be the submatrix obtained from  $A$  by deleting column  $C_1$

and the corresponding two rows in  $\Omega(C_1)$ . Then  $B$  is an inseparable  $(m-2) \times (n-1)$  orthogonal matrix of type 0. Therefore, we have

$$\omega(A) = \omega(B) + 8 \geq 2(m-2+n-1) - 2 + 8 = 2(m+n),$$

which is impossible. Therefore, we have shown that for any column  $C$  of weight 2 in  $A$ ,  $\tau(C) \geq 12$ . Similarly, for any row of weight 2 in  $A$ , the total number of 1's in the corresponding 2 columns is at least 12.

Let  $r$  (respectively,  $c$ ) be the number of rows (resp., columns) of weight 2 in  $A$ . Then by counting the total number of 1's in the columns of  $A$ , we have

$$\omega(A) \geq 12r + 2c + 4(n - 2r - c) = 4n + 4r - 2c.$$

Similarly, we have  $\omega(A) \geq 4m + 4c - 2r$ . This implies that

$$\omega(A) \geq 2(m+n) + r + c \geq 2(m+n),$$

which is a final contradiction. ■

## REFERENCES

1. L. B. Beasley, R. A. Brualdi, and B. L. Shader, Combinatorial orthogonality, in "Combinatorial and Graph-Theoretical Problems in Linear Algebra" (R. A. Brualdi, S. Friedland, and V. Klee, Eds.), pp. 207–218, Springer-Verlag, New York, 1993.
2. A. V. Geramita and J. Seberry, "Orthogonal Designs," Lecture Notes in Pure and Applied Mathematics, Vol. 45, Marcel Dekker, New York, 1979.
3. Peter M. Gibson and Guo-Hui Zhang, Combinatorially orthogonal matrices and related graphs, to appear in "Linear Algebra and its Applications".
4. K. B. Reid and C. Thomassen, Edge sets contained in circuits, *Israel J. Math.* **24** (1976), 305–319.
5. Guo-Hui Zhang, The spectrum of the number of 1's in a fully indecomposable orthogonal matrix over  $\mathbb{Z}_2$ , preprint.
6. Guo-Hui Zhang, Quasi-weighing matrices, preprint.